Pre & Post Care Guidelines

PRE-TREATMENT PREPARATIONS

- · Discontinue use of Retin A, Retinols, Vitamin A creams and other topic medications for 4 days before treatment.
- Daily Use of SPF 30+
- Tyrosinase inhibiters should be used for a minimum of 2 weeks prior to treatment for a Fitzpatrick 3 and higher.
- Series of 3 LED treatments is recommended to enhance results.

POST-TREATMENT

DO:

- Use an Image post treatment kit for the following 4-7 days
- · Avoid direct sun exposure for 2 weeks
- Use SPF 30+ or higher daily after 12 hours
- Wash the face a few hours after treatment or before bed time using a gentle cleanser and tepid water. This will improve the appearance of the skin and also allow for better subsequent absorption of products applied to the treated area
- 12-24 hours post-treatment mineral makeup can be applied
- LED Treatments immediately after with Blue (415nm) and Near Infrared (830nm))

DO NOT: Scrub, pick or rub your skin, as this can lead to scarring.

| AVOID | 24 HOURS | 2 DAYS | 4-7 DAYS | 7 DAYS | 7-14 DAYS |
|------------------------|----------|--------|----------|--------|-----------|
| Sunscreen | X | | | | |
| Mineral Makeup | X | | | | |
| Hot Shower | X | | | | |
| Shaving | X | | | | |
| Exercise | | X | | | |
| Spas & Sauna | | X | | | |
| Self Tanners | | X | | | |
| Swimming | | X | | | |
| AHAs & BHAs | | | X | | |
| Benzoyl Peroxide | | | X | | |
| Retinoids | | | X | | |
| Hydroquinone | | | X | | |
| Alcohol-Based Products | | | X | | |
| Wax/Sugar | | | | X | |
| Direct Sun Exposure | | | | | X |